



Section 24(c) Special Local Need Label

S-METOLACHLOR

GROUP 15 HERBICIDE

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF KANSAS

Dual MAGNUM®

EPA Reg. 100-816 **EPA SLN No. KS-100001**

KEEP OUT OF REACH OF CHILDREN

CAUTION

SYNGENTA'S SPECIAL CONDITIONS, RISKS OF USE AND DISCLAIMER FOR USE OF **DUAL MAGNUM ON CROPS ON THIS 24(C) LABEL**

IMPORTANT- READ BEFORE USE

THESE CONDITIONS RISKS OF USE AND DISCLAIMER ARE REQUIRED BY SYNGENTA CROP PROTECTION, INC AND NOT SPECIFIED BY U.S. EPA OR THE STATE OF KANSAS

Labeling may be obtained through the Special Labels link at www.farmassist.com.

FOR WEED CONTROL IN SESAME

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SYNGENTA CROP PROTECTION, INC. INTENDS THAT THE PRODUCT THAT IS THE SUBJECT OF THIS SECTION 24(C) LABEL BE PURCHASED ONLY BY END USERS WHO AGREE BY ELECTRONIC SIGNATURE ON SYNGENTA CROP PROTECTION'S INTERNET SITE TO THE TERMS AND CONDITIONS REQUIRED BY SYNGENTA CROP PROTECTION. INC. INCLUDING A WAIVER AND RELEASE FROM ALL LIABILITY AND INDEMNIFICATION BY THE USER AND/OR GROWER OF SYNGENTA AND OTHERS ARISING OUT OF OR RELATING TO ANY (1) FAILURE TO PERFORM, CROP INJURY, CROP YIELD REDUCTION, AND/OR CROP LOSS FROM USE OF DUAL MAGNUM ON CROPS ON THIS 24(C) LABEL; AND/OR (2) FAILURE TO COMPLY WITH THE TERMS AND CONDITIONS OF THIS 24(C) LABEL, INCLUDING THE RESTRICTIONS LISTED BELOW. IF SUCH TERMS AND CONDITIONS ARE UNACCEPTABLE, RETURN THE DUAL MAGNUM AT ONCE UNOPENED OR USE THE DUAL MAGNUM FOR A DIFFERENT APPROVED USE IN ACCORDANCE WITH THE LABEL AFFIXED TO THE PRODUCT CONTAINER.

USE OF DUAL MAGNUM (THE "PRODUCT") ON CROPS LISTED (THE "CROP") FOR THIS SPECIAL LOCAL NEED MAY RESULT IN CROP INJURY. CROP YIELD REDUCTION AND/OR CROP LOSS AS FURTHER DISCUSSED BELOW. READ AND UNDERSTAND THESE CONDITIONS AND RISKS OF USE FOR SPECIAL LOCAL NEED BEFORE USING

THE PRODUCT ON THE CROP. SYNGENTA RECOMMENDS THAT THE USER TEST THIS PRODUCT TO DETERMINE ITS SUITABILITY FOR SUCH INTENDED USE.

Syngenta Crop Protection, Inc. makes the Product available for use in the manner described in this Supplemental Labeling on the basis that, in the sole opinion of the user, the benefits and utility derived from the use of the Product on the Crop outweigh the potential risk of Crop injury, Crop yield reduction or Crop loss. The decision to use this Product in the manner described in this Supplemental Labeling must be made by each individual user on the basis of anticipated benefits versus (i) the potential risk of Crop injury, Crop yield reduction and Crop loss, (ii) the severity of the target pest infestation, (iii) the cost and availability of alternative pest controls and (iv) any other relevant factors. Syngenta recommends that the user test this Product to determine its suitability for such intended use.

By purchasing the Product for use, or using the Product in the manner described in this Supplemental Labeling, you acknowledge and accept that, to the extent consistent with applicable law:

- 1) you assume all risk of Crop injury, Crop yield reduction and Crop loss;
- 2) Syngenta Crop Protection, Inc. do not make, and do not authorize any agent or representative to make, any representations or recommendations regarding the use of this Product on the Crop other than the statements on this Supplemental labeling;
- 3) Syngenta Crop Protection, Inc. do not make, and do not authorize any agent or representative to make, any warranties, express or implied, with respect to the use of the Product on the Crop and disclaim all warranties, expressed or implied, including any implied warranty of merchantability:
- 4) Syngenta Crop Protection, Inc. disclaim all liability for any damages, losses, expenses, claims or causes of actions arising out of or relating to any (i) Crop injury, Crop yield reduction and/or Crop loss; and/or (ii) non-compliance with the terms and conditions of this Supplemental Labeling, including the restrictions listed below.
- 5) these conditions and Risks of Use for Special Local Need supersede any contrary representations or recommendations by Syngenta Crop Protection, Inc. or their respective agents or representatives, and any provisions in or on any Product literature or labeling including any provisions on the label affixed to the Product container.

If these Conditions and Risks of Use for Special Local Need are not acceptable, the unopened Product may be returned to the seller for a refund or used for a different labeled use in accordance with the label affixed to the Product container.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY OR LOSS, OR ILLEGAL RESIDUES.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR DISEASE CONTROL, CROP INJURY OR LOSS, OR ILLEGAL RESIDUES.

Follow all applicable directions, resistance management guidelines, restrictions, and precautions, including statements pertaining to the worker protection standards, on the Dual MAGNUM (EPA RegistrationNo.100-816) label.

This label must be in the possession of the user at the time of pesticide application.

SESAME

Dual MAGNUM can be used in sesame for pre-emergence suppression or control of annual grass and small-seeded broadleaf weeds. Dual MAGNUM will not control emerged weeds. Control emerged weeds with an appropriate registered foliar herbicide or by mechanical or physical means.

Attention: Read following "Restrictions" statement before proceeding.

Restrictions:

- 1. Do not make more than one application of Dual MAGNUM per crop per year.
- Do not apply more than 1.33 pts/A (1.27 lb ai/A) of Dual MAGNUM per crop per year. Do
 not apply more than 1.27 lb ai/A per year of S-metolachlor-containing products.
- 3. Harvest at normal maturity but no closer than 100 days following application.

Broadcast Application: Apply Dual MAGNUM as a broadcast, pre-emergence soil surface treatment at 0.67-1.33 pts/A (0.64 - 1.27 lb ai/A) after the crop has been planted but before the crop and weeds emerge. Use the lower end of the Dual MAGNUM rate range for soils that are relatively coarse textured (sandy soils) and the higher rates on fine textured soils (silt and clay soils).

Band Application: For sesame grown on beds or in rows, Dual MAGNUM may be applied as a pre-emergence (after planting but before crop and weed emergence) surface banded treatment over the row. For band applications, using row and band width measurements in inches, calculate the amount to be applied per acre as follows:

Band width in inches
Row width in inches

Dual MAGNUM Rate per acre for a broadcast treatment

Amount of Dual
MAGNUM needed per
acre for a banded
treatment

Precautions:

- 1) Dual MAGNUM will not control emerged weeds. Control emerged weeds with an appropriate registered foliar herbicide or by mechanical or physical means.
- 2) Do not irrigate after the Dual MAGNUM application and prior to sesame emergence or significant crop injury may occur.
- 3) If an activating rainfall (0.5") does not occur within 7-10 days following application, weed control may be reduced.
- 4) Do not incorporate Dual MAGNUM or severe sesame injury may occur.
- 5) Do not apply this product through any type of irrigation system
- 6) Excessive rainfall (> 1") following application will increase the risk for sesame injury.
- 7) Cool and wet weather will slow the growth of sesame and can increase the risk of crop injury.

Dual MAGNUM® is a trademark of a Syngenta Group Company

©2010 Syngenta Crop Protection

Section 24(c) Registrant: Syngenta Crop Protection, Inc. Greensboro, North Carolina 27419

KS0816026BA0910