

KANSAS DEPARTMENT OF AGRICULTURE**OFFICIAL****KUDZU CONTROL PROGRAM**

Revised January 1, 2004

DESCRIPTION

Kudzu, (*Pueraria lobata*), is a long-lived, coarse, viney legume that covers the ground with long runners. The leaflets are found in groups of three (5-20 cm. in length). The leaflets are 2-3 lobed and abruptly taper to a pointed tip. The stems have rough bark-like covering. Large flowers, 15-20 cm., lavender to purple, set on seeds sparingly, because of sparse blooming. The seed pods, 4.5 to 5 cm. in length are papery and covered with fine hair. The long runners root at the nodes to form new plants. Crowns taken from old stands are used for planting.

PREVENTION OF SPREAD OF KUDZU

The occurrence of new infestations of Kudzu can be reduced by preventing the movement of root crowns or seed from infested areas.

KUDZU CONTROL PRACTICES

Kudzu should be eradicated as quickly as possible by approved chemical. Control of Kudzu shall mean preventing the production of viable seed and destroying the plant's ability to reproduce by vegetative means.

CULTURAL CONTROL PRACTICES

Cultural control methods for Kudzu are not developed at this time.

HERBICIDES APPROVED FOR CONTROLLING KUDZU

The following herbicides may be used for cost-share with landowners. Other products labeled and registered for use on this noxious weed in Kansas may be used in accordance with label directions but are not available for cost-share.

Be sure to follow all label directions and precautions. For additional information consult the current KSU publication of AChemical Weed Control for Field Crops, Pastures, Rangeland, and Noncropland@.

Dicamba (Banvel, Clarity, Vanquish)

Glyphosate (Roundup)

Picloram (Tordon)

Triclopyr (Remedy, Garlon)

BIOLOGICAL CONTROL PRACTICES

There are no biological controls approved for Kudzu control at this time.